

NOW APPROVED FOR
**Mycoplasma
 bovis**

PREMIUM ANTIBIOTICS LABELED FOR THE TREATMENT OF BOVINE RESPIRATORY DISEASE

BRD PATHOGENS

ANTIBIOTIC	CONTROL OF BRD- ASSOCIATED PYREXIA (FEVER)	BRD PATHOGENS				BACTERICIDAL *
		MYCOPLASMA BOVIS	MANNHEIMIA HAEMOLYTICA	PASTEURILLA MULTOCIDA	HISTOPHILUS SOMNI	
Resflor Gold® (florfenicol)	X	X	X	X	X	X**
Nuflor Gold® (florfenicol)		X	X	X	X	X**
Draxxin® (tulathromycin)		X	X	X	X	
Baytril® (enrofloxacin)			X	X	X	X
Nuflor® (florfenicol)			X	X	X	X**
Excede® (ceftiofur)			X	X	X	X
Micotil® (tilmicosin)			X	X	X	

* An *in vitro* measure. The correlation between *in vitro* data and clinical effectiveness is unknown.
 ** Exhibits bactericidal activity against some strains of *Mannheimia haemolytica* and *Histophilus somni*.

See a Difference in One Dose





Florfenicol and Funixin Meglumine

Antimicrobial/Non-Steroidal Anti-inflammatory Drug

For subcutaneous use in beef and non-lactating dairy cattle only. Not for use in female dairy cattle 20 months of age or older or in calves to be processed for veal.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: RESLOR GOLD[®] is an injectable solution of the synthetic antibiotic florfenicol and the nonsteroidal anti-inflammatory drug (NSAID) funixin. Each milliliter of sterile RESLOR GOLD[®] contains 300 mg florfenicol, 6.5 mg funixin as funixin meglumine, 300 mg 2-glycine, 50 mg maleic acid, and triethanolin.

INDICATION: RESLOR GOLD[®] is indicated for treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis* and control of BRD-associated pyrexia in beef and non-lactating dairy cattle.

DOSE AND ADMINISTRATION: RESLOR GOLD[®] should be administered once by subcutaneous injection at a dose rate of 40 mg florfenicol/kg body weight and 2.7 mg funixin/kg body weight (6 mL/100 lb). Do not administer more than 10 mL at each site. The injection should be given only in the neck. Injection sites where the neck has not been evaluated for the 500mL, vial do not purchase the vial unless you are sure you will use it.

RESLOR GOLD [®] Dosage Guide ^a	
Animal Weight (lbs)	Dosage (mL)
100	6.0
200	12.0
300	18.0
400	24.0
500	30.0
600	36.0
700	42.0
800	48.0
900	54.0
1000	60.0

^a Do not administer more than 10 mL at each site.

CONTRAINDICATIONS: Do not use in animals that have shown hypersensitivity to florfenicol or funixin.

WARNINGS: NOT FOR HUMAN USE. KEEP OUT OF REACH OF CHILDREN. This product contains material that can be harmful if inhaled. Avoid contact with eyes, nose, mouth, and skin. Wash hands thoroughly after handling. Do not use if the product has been opened for 15 minutes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. Consult a physician if irritation persists. Accidental injection of this product may cause local irritation. Consult a physician immediately. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information.

For customer service or to obtain a copy of the MSDS, call 1-800-211-5675. For technical assistance or to report suspected adverse reactions, call 1-800-219-9286.

PRECAUTIONS: As a class, cyclo-oxygenase inhibition (NSAIDs) may be associated with gastrointestinal, renal, and hepatic toxicity. Sensitivity to drug-associated adverse events varies with the individual patient. Patients at greatest risk for adverse events are those that are dehydrated, on diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be carefully monitored. NSAIDs may inhibit the prostaglandins that maintain normal hemostatic function. Such anti-prostaglandin effects may result in clinically significant increases in bleeding or bruising or in bleeding or bruising that have not been previously associated with NSAID therapy. NSAIDs may also interact with other anti-thrombotic agents, such as RESLOR GOLD[®] with other anti-thrombotic drugs, such as NSAIDs or contrast agents, should be avoided or closely monitored.

Florfenicol is a cyclic oxazolidinone NSAID, and its effects in this class, adverse effects may occur with its use. The most frequently reported adverse effects have been gastrointestinal signs. Events involving reported renal, hematologic, neurologic, dermatologic, and hepatic effects have also been reported for other drugs in this class.

Not for use in animals intended for breeding purposes. The effects of florfenicol on bovine reproductive performance, pregnancy, and lactation have not been determined. Toxicity studies in dogs, rats, and mice have associated the use of florfenicol with testicular degeneration and atrophy. NSAIDs are known to have potential effects on both parturition and the estrous cycle. There may be a delay in the onset of estrus if funixin is administered during the proestrus phase of the estrous cycle. The effects of funixin on imminent parturition have not been evaluated in a controlled study. NSAIDs are known to have the potential to delay parturition through a "toxicity effect."

RESLOR GOLD[®] when administered as directed, may induce a transient reaction at the site of injection and underlying tissues that may result in trim loss of edible tissue at slaughter.

RESIDUE WARNINGS: Animals intended for human consumption must not be slaughtered within 38 days of treatment. Do not use in female dairy cattle 20 months of age or older. Use of florfenicol in this class of cattle may cause milk residues. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

ADVERSE REACTIONS: Transient hypotension, diarrhea, decreased water consumption, and injection site swelling have been reported. In some cases, the injection site swelling may be severe. In some cases, hypotension and collapse have been reported. Post-mortem examination of animals that died after treatment with RESLOR GOLD[®] has revealed the presence of florfenicol in cattle.

In cattle, rare instances of nephrotoxic-like reactions, some of which have been fatal, have been reported, primarily following intramuscular use of funixin meglumine.

CLINICAL PHARMACOLOGY: The pharmacokinetics (PK) of florfenicol (Table 1) and funixin (Table 2) after subcutaneous injection of RESLOR GOLD[®] is described below.

Table 1. Mean (n=28) pharmacokinetic parameters for florfenicol in cattle after a single subcutaneous administration of RESLOR GOLD (florfenicol dose of 40 mg/kg BW).

PK Parameter	Mean Florfenicol PK parameters in Cattle				
	AUC _{0-12h} ¹ (ng ² /h/mL)	AUC _{0-12h} ² (ng ² /h/mL)	C _{max} (ng/mL)	T _{1/2} (hr)	MRT _{0-12h} ³ (hr)
Mean	24527	247577	11151	6.25	28.5
SD	4274	41391	4194	3.97	11.6

Table 2. Mean (n=28) pharmacokinetic parameters for funixin in cattle after a single subcutaneous administration of RESLOR GOLD (funixin dose of 2.2 mg/kg BW).

PK Parameter	Mean Funixin PK parameters in Cattle				
	AUC _{0-12h} ¹ (ng ² /h/mL)	AUC _{0-12h} ² (ng ² /h/mL)	C _{max} (ng/mL)	T _{1/2} (hr)	MRT _{0-12h} ³ (hr)
Mean	13370	144488*	1913	1.14	9.5**
SD	4864	5116	791	0.97	3.27

¹ AUC_{0-12h} = Area under the plasma concentration-time curve (AUC) from time zero to the last quantifiable concentrations

² AUC_{0-12h} = AUC from time zero to infinity

³ C_{max} = Maximum plasma concentration

⁴ T_{1/2} = Terminal elimination half-life

⁵ MRT = Mean residence time from time zero to infinity

SD = Standard deviation

** p < 0.02

MICROBIOLOGY: Florfenicol is a synthetic, broad-spectrum antibiotic against many Gram-negative and Gram-positive bacteria isolated from domestic animals. It acts by binding to the 50S ribosomal subunit and inhibiting bacterial protein synthesis. Florfenicol is generally considered a bacteriostatic drug, but exhibits bactericidal activity against certain bacterial species. *In vitro* studies demonstrate that florfenicol is active against the BRD pathogens *M. haemolytica*, *P. multocida*, *H. somni* and *M. bovis* and that florfenicol exhibits bactericidal activity against strains of *M. haemolytica* and *H. somni*.

The minimum inhibitory concentrations (MICs) of florfenicol were determined for BRD isolates obtained from calves enrolled in BRD field studies in the U.S. in 2006 using methods recommended by the Clinical and Laboratory Standards Institute (M7-A2). Isolates were obtained from pre-treatment nasal swabs from all calves enrolled at all four sites, post-treatment nasal swabs from treatment failures in the RESLOR GOLD and saline control treatment groups at three sites, and lung tissue from one calf that died in the saline control treatment group. The results are shown in Table 3.

Table 3. Florfenicol MIC values^a of indicated pathogens isolated from cattle with naturally-occurring BRD.

Indicated pathogens	Year of isolation	Number of isolates	MIC ₅₀ ** (µg/mL)	MIC ₉₀ ** (µg/mL)	MIC range (µg/mL)
<i>Mannheimia haemolytica</i>	2006	183	1.0	1.0	0.5 to 32
<i>Pasteurella multocida</i>	2006	139	0.5	0.5	≤ 0.125 to 16
<i>Histophilus somni</i>	2006	84	≤ 0.125	≤ 0.125	≤ 0.125 to 0.25

^a The correlation between *in vitro* susceptibility data and clinical effectiveness is unknown.

^b The lowest MIC to encompass 50% and 90% of the most susceptible isolates, respectively.

EFFECTIVENESS: In a multi-site field study, calves with naturally-occurring BRD were treated with RESLOR GOLD[®], Nullor GOLD[®] (Mylan, NJ, 286), or saline. A treatment success was defined as a return to normal respiration to mild respiratory signs, normal attitudes to touch, and a rectal temperature < 104.0 °F on Day 11.

The treatment success rate for BRD for the RESLOR GOLD[®] treatment group (68.8%) was statistically significantly greater ($p < 0.0002$) compared to control treatment groups (39.9% for Nullor GOLD[®] and 39.9% for saline). The difference in the treatment of BRD, with one-sided 85% lower confidence bound for the difference between the two treatment groups $p < 0.23\%$.

In the same study, the change in rectal temperature from pre-treatment to six hours post-treatment was evaluated to assess the effect of treatment on fever. The mean rectal temperature at six hours post-treatment was statistically significantly greater ($p < 0.0019$) in the RESLOR GOLD[®] treatment group compared to the saline control treatment group. The mean rectal temperatures decreased by > 2.0 °F from pre-treatment to six hours post-treatment was statistically significantly greater in rectal temperature from pre-treatment to six hours post-treatment was statistically significantly greater in the RESLOR GOLD[®] treatment group compared to the Nullor GOLD[®] and saline control treatment groups ($p < 0.031$ and 0.002, respectively).

The effectiveness of RESLOR GOLD for the treatment of BRD associated with *Mycoplasma bovis* was demonstrated by examining the *M. bovis* data from cattle enrolled in the BRD treatment failures (2 of 8 calves, 25%) in RESLOR GOLD-treated calves that cultured positive for *M. bovis* post-treatment.

ANIMAL SAFETY: A target animal safety study was conducted to evaluate the effects of RESLOR GOLD[®] when administered to cattle subcutaneously at 1X, 3X, or 5X the label dose for five consecutive days (D5) in the United States. Decreased feed and water consumption were observed in the 1X, 3X, and 5X groups. Injection site swellings were observed in the 1X, 3X, and 5X groups.

A separate injection site study was conducted in cattle. The study demonstrated that RESLOR GOLD[®], when administered according to the label directions, may induce a transient reaction in the subcutaneous and underlying muscle tissue.

STORAGE INFORMATION: Do not store above 30 °C (86 °F). Use within 28 days of first use.

HOW SUPPLIED: RESLOR GOLD[®] is available in 100, 250, and 500 mL sterile, multiple-dose, glass vials.